Removal of Radial Hemostatic Band (Terumo TR Band) Competency Verification Record University of Virginia Health System

Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	Employee Name:	Employee ID #:	Date:	
Instructions: Successful completion is documented on the Annual Competency Record (ACR): At the end of training session the participant will be able to: Demonstrates procedure to remove TR band. Demonstrated Skill: Gathers Supplies: Goggles/Face Mask or Face Shield TR Band Syringe (Only for TR Band) Pulse oximeter (if available, to assist with perfusion monitoring) Sterile 2x2 Clear transparent dressing Penlight or other light source (to assist with site assessment for bleeding during removal) Arm board Dons PPE. Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	as a guide for competency check of competency validation occurs awa be presented to the unit NEC or m	ff only; the Annual Competency Record is used to by from the unit, this form can be completed by th anager as evidence of competency. The Annual C	o document competen e validator; the signe	cy. (If d form can then
Successful completion is documented on the Annual Competency Record (ACR): At the end of training session the participant will be able to: Demonstrates procedure to remove TR band. Demonstrated Skill: Beautiful Scatter Supplies: Gloves Goggles/Face Mask or Face Shield TR Band Syringe (Only for TR Band) Pulse oximeter (if available, to assist with perfusion monitoring) Sterile 2x2 Clear transparent dressing Penlight or other light source (to assist with site assessment for bleeding during removal) Arm board Dons PPE. Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes to wait 1 minute after all the air is removed (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work		s vanaaiea).		
Gathers Supplies: Gloves Goggles/Face Mask or Face Shield TR Band Syringe (Only for TR Band) Pulse oximeter (if available, to assist with perfusion monitoring) Sterile 2x2 Clear transparent dressing Penlight or other light source (to assist with site assessment for bleeding during removal) Arm board Dons PPE. Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Competency Verified by:	 Successful completion is docume At the end of training session the 	participant will be able to:		
Gathers Supplies: Gloves Gloves Gorgeles/Face Mask or Face Shield TR Band Syringe (Only for TR Band) Pulse oximeter (if available, to assist with perfusion monitoring) Sterile 2x2 Clear transparent dressing Penlight or other light source (to assist with site assessment for bleeding during removal) Arm board Dons PPE. Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work	Demonstrates procedure to remove	ve TR band.		
Gloves Goggles/Face Mask or Face Shield TR Band Syringe (Only for TR Band) Pulse oximeter (if available, to assist with perfusion monitoring) Sterile 2x2 Clear transparent dressing Penlight or other light source (to assist with site assessment for bleeding during removal) Arm board Dons PPE. Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	I	Demonstrated Skill:		
Gloves Goggles/Face Mask or Face Shield TR Band Syringe (Only for TR Band) Pulse oximeter (if available, to assist with perfusion monitoring) Sterile 2x2 Clear transparent dressing Penlight or other light source (to assist with site assessment for bleeding during removal) Arm board Dons PPE. Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	Gathers Supplies:			
 TR Band Syringe (Only for TR Band) Pulse oximeter (if available, to assist with perfusion monitoring) Sterile 2x2 Clear transparent dressing Penlight or other light source (to assist with site assessment for bleeding during removal) Arm board Dons PPE. Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by: 				
 Pulse oximeter (if available, to assist with perfusion monitoring) Sterile 2x2 Clear transparent dressing Penlight or other light source (to assist with site assessment for bleeding during removal) Arm board Dons PPE. Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by: 	 Goggles/Face Mask or Face S 	hield		
 Sterile 2x2 Clear transparent dressing Penlight or other light source (to assist with site assessment for bleeding during removal) Arm board Arm board Dons PPE. Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by: 	• TR Band Syringe (Only for T	R Band)		
Clear transparent dressing Penlight or other light source (to assist with site assessment for bleeding during removal) Arm board Dons PPE. Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	 Pulse oximeter (if available, to 	o assist with perfusion monitoring)		
Penlight or other light source (to assist with site assessment for bleeding during removal) Arm board Dons PPE. Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	• Sterile 2x2			
• Arm board Dons PPE. Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	ž			
Dons PPE. Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:		(to assist with site assessment for bleeding d	luring removal)	
Verbalizes monitoring required while band is in place (q15 min site check and perfusion monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:				
monitoring using O2 sat probe on thumb or index finger distal to band; if O2 sat monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:				
monitoring not in place, monitor cap refill, skin color/temp). Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:				
Verbalizes timing of removal. Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:			02 sat	
Elevate the patient's arm on 1-2 pillows or place a pillow upward on the bedrail so that the wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	Ţ .	p refill, skin color/temp).		
wrist is higher than the elbow. Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:				
Attach syringe and slowly removes 3-5 ml of air q5 minutes until all air has been removed. Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:		lows or place a pillow upward on the be	drail so that the	
Removes the syringe from the port after each air removal. Purges all except 1 ml of air from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	3			
from the syringe before next air removal. Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	· ·			
Re-inject 1-2 ml of air or the amount of air it takes to stop a re-bleed. (Wait 15-30 minutes and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:			t 1 ml of air	
and repeat the steps above.) Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:				
Verbalizes to wait 1 minute after all the air is removed, then carefully removes the Radial Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	Re-inject 1-2 ml of air or the amoun	nt of air it takes to stop a re-bleed. (Wait	t 15-30 minutes	
Hemostatic Band from the wrist. Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	and repeat the steps above.)			
Places a sterile 2X2 and clear transparent dressing over the access site. Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	Verbalizes to wait 1 minute after al	I the air is removed, then carefully remo	ves the Radial	
Verbalizes action to take is a re-bleed occurs after band removal (manual pressure). Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:				
Places arm board to back of forearm and secures. (Verbalizes this is to remain on for at least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	Places a sterile 2X2 and clear transp	parent dressing over the access site.		
least 6 hrs and up to 24 hours, do not cover site/dressing.) Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	Verbalizes action to take is a re-ble	ed occurs after band removal (manual pr	ressure).	
Verbalizes EPIC documentation process and where to find resources for documentation and standard work Competency Verified by:	Places arm board to back of forearr	n and secures. (Verbalizes this is to rema	ain on for at	
and standard work Competency Verified by:				
Competency Verified by:		ocess and where to find resources for do	cumentation	
	and standard work			
Data	Competency Verified by:		Date:	

Evaluator's signature

Evaluator's Name (printed)